US ERA ARCHIVE DOCUMENT

ASSESSING THE QUALITY AND QUANTITY OF DEPRESSIONAL WETLANDS IN THE REDWOOD RIVER WATERSHED UTILIZING A PROBABILISTIC SURVEY DESIGN

John Genet

MN Pollution Control Agency-St. Paul, MN


Anthony R. Olsen
US EPA NHEERL-Corvallis, OR

EMAP 2007 Symposium April 10-11th 2007 Washington, DC

Introduction

Index of Biological Integrity (IBI)

- Multimetric Index
- Based on deviation from reference conditions (least impacted sites)
- Measurement of condition (quality)






Introduction

Assessment

- Aquatic LifeUse Support(ALUS)
- Impairment threshold
 - Lowest scoring reference site
- Confidence interval






Study Objectives

- Assess quality of depressional wetlands in a watershed using a probabilistic survey
- Estimate wetland loss since early 1980s
- Evaluate feasibility of utilizing a randomized survey design for wetland condition monitoring



Study Area-Redwood River





Target Population

Seasonal, Semi-permanent, & Permanent Emergent Depressional Wetlands











Sample Frame-NWI

Redwood River Watershed Sample Frame: Depressional Wetlands




Site Selection & Sampling




Results-Water Chemistry

Chloride



Total N




--- Regional Reference Conditions




Results-Invertebrate IBI

Number of Basins

Cumulative Area





-- - Biological Impai

Biological Impairment Threshold

<u>Impairment</u>


Basins - 86%

Area - 97%




Results-Plant IBI

Number of Basins



Cumulative Area



Biological Impairment Threshold


<u>Impairment</u>

Basins – **63%**

Area - 81%




Results-ALUS




Error bars = 95% Confidence limits



Results-ALUS




Results-Site Evaluation






Results-Basin Extrapolation





Results-Area Extrapolation





Results-Wetland Losses

Size	Estimated No. of Wetland Basins			Estimated Wetland Area (ha)		
Category (ha)	circa 1980	2003	% change	circa 1980	2003	% change
< 1	1475	536	-63.6	486.8	201.7	-58.6
1-5	407	241	-40.9	848.2	541.2	-36.2
5-10	58	39	-33.3	396.2	266.8	-32.7
> 10	51	48	-6.2	1741.7	1711.1	-1.8
Total	2012	896	-55.5	3437.6	2714.4	-21.0




Conclusions

- Majority of depressional wetlands in the Redwood are biologically impaired
- Likely stressors: nutrients, invasive species, sedimentation, altered hydrology
- Significant losses of small (seasonal) wetland basins from 1980-2003
- These are not net losses
 - No accounting for permitted actions & mitigation, and/or wetland additions from conservation programs (CREP, WRP, etc)



Conclusions-Probabilistic Survey

- Accurate sample frame is crucial
 - NWI format is problematic
 - Data is > 20 yrs
- Access not a big issue





Acknowledgements

<u>Mark Gernes</u> – idea for project, study design, field work, comments & suggestions on report

Michael Bourdaghs - comments & suggestions on report, assistance with presentation

Field Crew

Joel Chirhart

Harold Wiegner

Maureen Minister

Bruce Sandstrom

Adam Hoffman

Travis Fristed

EPA Support

Rich Sumner

Mary Kentula

Naomi Detenbeck


Sue Elston

Alicia Hernandez



Comprehensive Wetland Assessment, Monitoring, and Mapping Strategy (CWAMMS)

- Probabilistic Survey of Wetland Quantity and Quality (Statewide Status & Trends)
- Update NWI
- Develop Web-Based Wetland Accounting System (permit and mitigation tracking)




Probabilistic Statewide Wetland Survey

3-Panel Interpenetrating Design








Year (Panel) 2 1580 1 mi² plots




Year (Panel) 3 1580 1 mi² plots

+ 250 Common Plots Sampled each year ----->

Panel 1: 2006 Wetland Quantity Status 2007 Wetland Quality Status





Questions?



Redwood Report available at:

http://www.pca.state.mn.us/water/biomonitoring/bio-wetlands.html

More CWAMMS information available at:

http://www.pca.state.mn.us/water/wetlands/cwamms.html